

The 2003 Superoutburst of an SU UMa-type Dwarf Nova, GO Comae Berenices

Akira IMADA¹, Taichi KATO¹, Makoto UEMURA¹, Ryoko ISHIOKA¹, Thomas KRAJCI²
Yasuo SANO³, Tonny VANMUNSTER⁴, Donn R. STARKEY⁵, Lewis M. COOK⁶, Jochen PIETZ⁷
Daisaku NOGAMI⁸, Bill YEUNG⁹, Kazuhiro Nakajima¹⁰, Kenji TANABE¹¹, Mitsuo KOIZUMI¹¹
Hiroki TAGUCHI¹¹, Norimi YAMADA¹¹, Yuichi NISHI¹¹, Brian MARTIN¹², Ken'ichi TORII¹³
Kenzo KINUGASA¹⁴ and Christopher P. JONES¹⁵

¹Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502
a.imada@kusastro.kyoto-u.ac.jp

²3933 Stockton Loop SE Albuquerque, New Mexico 87118-1104, USA

³VSOLJ, 3-1-5 Nishi-juni-jou minami, Nayoro, Hokkaido, Japan

⁴Center for Backyard Astrophysics (Belgium), Walhostraat 1A, B-3401, Landen, Belgium

⁵AAVSO, 2507 County Road 60, Auburn, Auburn, Indiana 46706, USA

⁶Center for Backyard Astrophysics (Concord), 1730 Herix Ct. Concord, California 94518, USA

⁷Rostocker Str. 62, 50374 Erftstadt, Germany

⁸Hida Observatory, Kyoto University, Kamidakara, Gifu 506-1314

⁹PO Box 105 Benson, Arizona 85602, USA

¹⁰VSOLJ, 124 Isatotoyo Teradani, Kumano, Mie, Japan

¹¹Department of Biosphere-Geosphere Systems, Faculty of Informatics, Okayama University of Science,
Ridaicho 1-1, Okayama 700-0005

¹²King's University College, Department of Physics, 9125 50th Street, Edmonton, AB T5H 2M1, Canada

¹³Department of Earth and Space Science, Graduate School of Science, 1-1 Machikaneyama-cho,
Toyonaka, Osaka 560-0043 Japan

¹⁴Gunma Astronomical Observatory, 6860-86 Nakayama Takayama-mura, Agatsuma-gun, Gunma 377-0702

¹⁵BAA, Variable Star Section, 29 Buller Road, Laindon, Essex SS15 6BA, England

(Received ; accepted)

Abstract

We photometrically observed the 2003 June superoutburst of GO Gom. The mean superhump period was 0.063059(13) d. The resultant data revealed that (1) the obtained light curve contained a precursor, (2) a plateau stage of the object lasted 8 days, which is remarkably shorter than that of ordinary SU UMa-type dwarf novae, and (3) the amplitude of the superoutburst was less than 5 mag, which is unpredictably small when taking into account the fact that the supercycle of GO Com is about 2800 days. In order to explain these anomalies, a mass elimination process from the accretion disk, such as evaporation, may play an important role.

Key words: accretion, accretion disks — stars: dwarf novae — stars: individual (GO Comae Berenices) — stars: novae, cataclysmic variables — stars: oscillations

1. Introduction

Dwarf novae are a class of cataclysmic variables (CVs) which are close binary systems consisting of a white dwarf and a late-type main-sequence secondary transferring gaseous matter via the Roche lobe overflow. The transferred gas forms an accretion disk around the white dwarf. The accretion disk gives rise to various kinds of instabilities and variabilities (for a review, see e.g., Warner 1995, Hellier 2001). The two major instabilities are thermal and tidal instabilities, which are responsible for dwarf nova-type outbursts and superhumps, respectively (for a review, see Osaki 1996). SU UMa-type stars, subclass of dwarf novae, satisfy both instabilities.

It is widely recognized that SU UMa-type dwarf novae have a subgroup, called WZ Sge-type dwarf no-

vae [originally proposed by Bailey (1979); see also Downes, Margon (1981), O'Donoghue et al. (1991), Kato et al. (2001b)]. WZ Sge-type dwarf novae are one of the most enigmatic systems all over the dwarf novae (see e.g., Osaki 1995, Lasota et al. 1995, Mineshige et al. 1998). There are only five objects that have been firmly classified into WZ Sge-type dwarf novae, HV Vir, AL Com, EG Cnc, RZ Leo, and WZ Sge itself. Recent paper (Kato et al. 2001b) lists possible candidates for WZ Sge-type dwarf novae. The mass ratios of these systems have been suggested to be much smaller than those of SU UMa-type dwarf novae.¹ Criteria for WZ Sge-type dwarf novae are that (1) a supercycle, which is a period between two suc-

¹ RZ Leo, however, is regarded as a system with a rather high mass ratio in comparison with the other WZ Sge-type dwarf novae.

ceeding superoutbursts, is over 5 years, much longer than those of SU UMa-type dwarf novae, (2) their outbursts show very large amplitudes exceeding 6 mag, and (3) near the maximum of a superoutburst, early superhumps, having doubly-peaked shapes and periods close to the P_{orb} , are observed for about a few days before emergence of ordinary superhumps.² Historically, Vogt, Bateson (1982) pointed out that GO Com is a candidate for WZ Sge-type dwarf novae.³ However, the lack of observations has prevented us from clarification.

GO Com was discovered by Kowal (1973) as an eruptive object on a Palomar plate taken on 1973 July 1.213. This variable was confirmed to be identical with the suspected variable star CSVS 1959 = SVS 382 discovered by Belyavskij (1933). Howell et al. (1990) derived the orbital period of GO Com to be 95 min.

Being located extremely close to the north galactic pole and in the SA 57 region, this object was independently recorded as a very blue object of $B=18.1$ (US 31) during a systematic survey of this region (Usher 1981). According to Usher (1981), this object has one of the most unusual colors ($U-B = -1.5$) in this region. This object is also cataloged as a quasar candidate OMHR J125637.12+263644.2 (Moreau, Reboul 1995).

The object has been visually monitored by the members of the Variable Star Observers League in Japan (VSOLJ) since 1986, and only one short outburst was observed on 1989 May 30 ($m_v=13.2$) until 1995. This 1989 outburst showed a rapid decline by about 1 mag d^{-1} , suggesting that the outburst would be a normal outburst of a short-period dwarf nova. Although there were unavoidable seasonal gaps in monitoring, the extremely low outburst frequency of this dwarf nova has been established by this observation. The apparent absence of long outbursts, infrequent short outbursts, and possible excursion to a low state (as indirectly inferred from the literature) have suggested similarity to the unusual short-period dwarf nova BZ UMa (Wenzel 1982; Kaluzny 1986; Ringwald, Thorstensen 1990; Jurcevic et al. 1994; Kato 1999; Neustroev et al. 2002). The relatively strong X-ray emission detected by ROSAT (Verbunt et al. 1997) also suggests the unusual nature of GO Com.

In 1990, Kato, Hirata (1990) conducted first-ever time-resolved CCD photometry of GO Com near its quiescence, and reported the possible presence of a periodicity of 33 min. Howell et al. (1990) found a possible 95-min periodicity by CCD photometry. Although this period was tentatively related to the possible orbital period, the short observation was not sufficient to conclude on the reality of this periodicity. We have had to wait for more than a decade before the true periodicity is confirmed by the present observation.

The 1995 July outburst was detected by Vanmunster on

1995 July 16, which was confirmed by CCD observation on July 25 at $V=16.6$ (Kato et al. 1995). The most unusual feature of this outburst was that this outburst preceded a long outburst by 14 d. This ‘second’ outburst very much looked like a superoutburst of SU UMa-type dwarf novae (Kato et al. 1995), although the limited visibility hindered secure detection of superhumps. The object even underwent a rebrightening following this outburst. All of these observations suggested that GO Com is a genuine SU UMa-type dwarf novae, and that the 1995 July outburst can be best interpreted as a precursor outburst to the following presumable superoutburst (cf. Marino, Walker 1979; Kato 1997).

Since then, there have been only four outbursts on the VSNET⁴ record: 1996 March, 1997 February, 1998 April, and 2003 June. The outbursts other than the last one were short and faint outbursts.

2. Observation

The CCD observations were performed by the international VSNET (Kato et al. 2004) collaboration team at 14 sites. A summary of the observations and the equipment used in each site is listed in table 1.

The images were dark-subtracted, flat-fielded and analyzed with the aperture photometry. The Kyoto and RIKEN images were analysed using a Java-based aperture and PSF photometry package developed by one of the authors (TK). The Hida images were analysed using IRAF. The other images were analyzed using AIP4WIN.

The magnitude scales of each observatory were adjusted to that of the most abundant Tashkent system (TKr in table 1). The magnitudes of the object were determined relatively using a local standard star, GSC1995.488, whose constancy during the run was confirmed using a check star, GSC1995.1151.

Heliocentric corrections to the observation times were applied before the following analysis.

3. Results

3.1. Long-term Monitoring of GO Com

As mentioned above, extensive monitoring of the object has been performed by many amateur astronomers. Data points which include negative observations amount to more than 4000 since 1995. However, despite the secure observations, only 2 superoutbursts have been detected. Table 2 represents the records of outbursts of GO Com. One can roughly estimate a supercycle of GO Com about 2800 days, judging from the two succeeding superoutbursts in 1995 July and 2003 June.

3.2. Light Curve

The resultant light curve of the superoutburst is shown in figure 1. We derived the decline rate of 0.15 mag d^{-1} during the plateau stage, which lasted for 8 days, from HJD2452796 to HJD2452804. The most significant fact

² The early superhump is sometimes called orbital hump (Patterson et al. 2002) or early hump (Osaki, Meyer 2002). The difference between these authors is originated from their interpretation of physical processes near the bright maximum

³ Note that criteria for TOADs in Howell et al. (1995a) are different from those for WZ Sge-type dwarf novae

⁴ <http://www.kusastro.kyoto-u.ac.jp/vsnet/>

is that the duration of the plateau stage is considerably shorter than that of normal SU UMa-type dwarf novae, whose duration is typically about 2 weeks (Table 3.7. in Warner (1995)).

After the termination of the steep decline, the object entered the post-superoutburst stage during which the object stayed more than about 1 mag brighter than in quiescence ($B=18.1$, mentioned above, and typically $B-V \sim 0$ in dwarf novae). A rebrightening feature is sometimes observed during this stage, especially among SU UMa-type dwarf novae with shorter orbital periods and WZ Sge-type dwarf novae. There is a hint of a rebrightening feature around HJD2452811, however, the lack of later data has hindered us from confirmation of rebrightening.

3.3. Superhumps

The result of a period analysis is depicted in figure 2, applying the PDM method (Stellingwerf 1978) to a data set covering the period from HJD2452795 to HJD2452801 after subtracting the linear decline trend. We obtained the mean superhump period P_{SH} of 0.06297(3) d. Figure 3 represents the phase-averaged mean superhump profile. The rapid-rise and slow-decline trend is characteristic of superhumps in SU UMa-type dwarf novae. This superhump period seems to be shorter than the orbital period derived by Howell et al. (1990). Precise measurement of the orbital period of GO Com is urged.

In order to investigate variations of superhump profile, we also derived the daily-averaged profiles of superhumps, shown in figure 4. The superhump amplitude on HJD2452795, corresponding to the very day when the superoutburst started, was as large as 0.4 mag, which gradually decayed during the following 8 days. However, at the end of a plateau stage, superhumps clearly regrew. Similar phenomena were observed SU UMa-type dwarf novae with longer supercycle (see e.g., Baba et al. 2000).

3.4. Superhump Period Change

We derived the timings of the superhump maxima mainly by eye. The accuracy of the eye estimates of the timings is around 0.001 d. The results are listed in table 3. A linear regression to the times yielded the following equation,

$$HJD(max) = 2452795.1515(15) + 0.063059(13) \times E(1)$$

where E is the cycle count since HJD2452795.1571. By fitting the deviations of the observed timings from the calculation, we obtained the following equation,

$$O - C = 0.0079(15) - 0.0005(1) \times E + 5.79(88) \times 10^{-6} \times E^2(2)$$

The quadratic term corresponds to $\dot{P}/P = 1.8 \times 10^{-4}$ (see figure 5). This means that the superhump period became longer, as shown in some SU UMa-type dwarf novae with shorter orbital periods (Kato et al. 2001b).

Figure 5 also gives us a hint of a late superhump, around HJD 2452802. However, it is difficult to examine it because of lacking of data.

3.5. Precursor Outburst

SU UMa-type dwarf novae sometimes undergo a superoutburst following a normal outburst (Marino, Walker 1979). We call such a normal outburst precursor. Thanks to the excellent coverage, the early stage of the present outburst is considered as a precursor before the superoutburst. Figure 1 indicates a decline of the precursor on HJD2452794.

Our calculation revealed that the mean decline rate of magnitude during the precursor was 0.74 mag d^{-1} . Figure 6 represents the precursor profile folded by $P=0.063059$ d after subtracting the decline trend. Judging from figure 6, there exists no superhump profile as is seen in a superoutburst of SU UMa-type dwarf novae.

4. Discussion

4.1. The nature of GO Com

As mentioned above, during a superoutburst, the amplitude and plateau stage of ordinary SU UMa-type dwarf novae are 4~6 mag and ~2 weeks, respectively (Warner 1995). On the other hand, during the 2003 superoutburst of GO Com, the amplitude and plateau stage of the object are ~5 mag and ~8 days, respectively. Especially, the plateau stage of GO Com is remarkably shorter than that of typical SU UMa-type dwarf novae. This may indicate that the accumulated mass on the accretion disk in quiescence is low in GO Com.

According to the disk instability theory, if an outburst is triggered in an accretion disk with a relatively low mass, the system is expected to have a short supercycle. However, in the case of GO Com, the observations suggest a long supercycle of ~2800 days and a moderately short superhump period of 0.063059 days. These results are beyond the framework of the standard disk instability model.

In order to account for the observations, we propose that high efficiency of mass accretion onto the white dwarf is realized during quiescence. This scenario was originally suggested by Lasota et al. (1995): the short plateau stage and the long supercycle are due to inner truncation of the accretion disk. Similar interpretation is discussed on LL And by Kato (2004). LL And is an SU UMa-type dwarf nova with an extremely long supercycle (~5000 d), comparable to that of WZ Sge-type dwarf novae. The obtained light curve of the 1993 superoutburst of LL And is remarkably similar to that of GO Com in terms of a small superoutburst amplitude and a short plateau stage. Table 4 demonstrates fundamental characteristics of GO Com and LL And. In conjunction with our observations, we conclude that the inner part of the accretion disk may be truncated, as modeled by Lasota et al. (1995).

Note, however, that various durations of superoutbursts have been observed in certain objects (e.g. SW UMa, Howell et al. (1995b)). Further observations are needed in order to clarify superoutburst properties of GO Com.

4.2. Superhump period change

In general, the period of superhump in SU UMa-type dwarf novae decreases during a superoutburst (e.g. Warner 1985; Patterson et al. 1993). The obtained $\dot{P}/P$ has a rather common negative value ($\sim 5 \times 10^{-5}$) (Kato et al. 2003b), which has been generally ascribed to decreasing apsidal motion due to a decreasing disk radius. Osaki (1985) pointed out mass depletion from the disk results in the decrease of superhump period.

However, recent observations reveal that the minority of SU UMa-type dwarf novae has positive $\dot{P}/P$ (Semeniuk et al. 1997; Nogami et al. 1998; Baba et al. 2000; Kato et al. 2001a; Kato, Starkey 2002; Olech 2003; Olech et al. 2003; Nogami et al. 2004).⁵ Kato et al. (2001b) suggested that an increase in superhump periods may be related to a low mass ratio and/or a low mass transfer rate. EI Psc (Uemura et al. 2002a), however, obviously violates these regime: Uemura et al. (2002b) derived a high mass ratio ($q = M_2/M_1 = 0.19 \pm 0.02$) by using the empirical relation between superhump excess and mass ratio (Patterson 2001). Spectroscopic observations for EI Psc also indicate the high mass transfer rate (Hu et al. 1998; Wei et al. 2001). It is suggested that another condition may be required for explaining positive $\dot{P}$.

Figure 7 demonstrates a $\dot{P}/P$ diagram of SU UMa-type dwarf novae (including WZ Sge-type dwarf novae). Note that the obtained $\dot{P}/P$ of GO Com is the second largest ever known. In addition, the location of GO Com in figure 7 are far from the general trend from upper-left to lower-right, implying a peculiar nature of GO Com.

4.3. Precursor and before the maximum

Observations before the maximum allow us to test the validity of a refinement of the thermal-tidal instability model developed by Osaki, Meyer (2003). Osaki, Meyer (2003) suggested that a superoutburst which is accompanied with a precursor should develop superhumps after the termination of the precursor, i.e., as soon as a superoutburst is triggered, superhumps emerge. In this point, our observations are in agreement with the prediction by Osaki, Meyer (2003).

We also investigated the possibilities of the presence of superhumps during the precursor. Kato (1997) found a hump-like profile with an amplitude smaller than 0.1 mag during a precursor of the 1993 superoutburst in T Leo. He suggested that this modulation and the superhump were caused by the same origin, based on the fact that the ephemeris of the superhump-maximum timings during the superoutburst was applicable to that of the modulations during the precursor. This implies that a tidal instability developed before the main superoutburst. However, in the case of GO Com, there is no evidence of superhumps during the precursor. The reason for the difference is left as an open question.

⁵ All WZ Sge-type dwarf novae show positive $\dot{P}$.

4.4. Relation to WZ Sge-type dwarf novae

As mentioned above, historical records indicate that the supercycle of GO Com is about 2800 day, which is considerably longer than well-known SU UMa-type dwarf novae (200-350 d) (Nogami et al. 1997, Kato et al. 2003a). Thus, from the point of supercycle, GO Com is similar to WZ Sge-type dwarf novae, although their supercycles are much longer than that of GO Com.

However, GO Com underwent normal outbursts more frequently than superoutbursts. Observations reveal that WZ Sge-type dwarf novae enter a normal outburst very infrequently.⁶ Theories predict that an extremely low viscosity on an accretion disk during quiescence leads WZ Sge-type dwarf novae to have a long dormancy (Osaki 1995). In the case of GO Com, assuming the analogy with WZ Sge-type dwarf novae, rather frequent normal outbursts may not be explained. Thus, the viscosity in the accretion disk is not necessarily low during quiescence.

We also investigated the possibility of the existence of an early superhump on HJD 2452795, one day before the maximum. However, the profiles we detected was not doubly peaked, but singly peaked. Therefore, the modulations on HJD 2452795 were definitely ordinary superhumps. In addition, precursor outbursts are not shown among WZ Sge-type dwarf novae. That's why we disqualify GO Com as WZ Sge-type dwarf novae.

5. Summary

Our photometrical study in this paper is summarized as follows.

1. GO Com was proved to be an SU UMa-type dwarf nova with superhumps whose period is specified as 0.063059 d.
2. The plateau stage of the superoutburst continued only 8 days, during which the period of superhump became longer.
3. A precursor was followed by the superoutburst. Our detailed analysis finds no evidence for superhumps during the precursor, which is in contrast with the results of Kato (1997) about T Leo.
4. Judging from long-term monitoring, the supercycle of GO Com is about 2800 d, whereas the amplitude of the 2003 superoutburst was less than 5 mag. In conjunction with its plateau stage of GO Com (8 days), the total mass accretion onto the white dwarf might be low.

We are grateful to many VSNET observers who have reported vital observations. We are particularly grateful to Christopher Jones for promptly reporting the outbursts through VSNET. This work is supported by the Grant-in-Aid for the 21st Century COE "Center for Diversity and Universality in Physics" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. This work is partly supported by a grant-in aid

⁶ There is no record of a normal outburst for WZ Sge itself.

[13640239 (TK), 14740131 (HY), 16740121 (KK)] from the Ministry of Education, Culture, Sports, Science and Technology. Part of this work is supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (MU, RI). This research has made use of the Digitized Sky Survey produced by STScI, the ESO Skycat tool, and the VizieR catalogue access tool.

Table 3. Timing of superhumps maxima during the super-outburst

HJD-2400000	E	$O - C$
52795.1571	0	0.0055
52795.2184	1	0.0038
52795.2807	2	0.0030
52795.7230	9	0.0039
52795.7843	10	0.0021
52796.0395	14	0.0051
52796.0995	15	0.0020
52796.1608	16	0.0003
52796.4147	20	0.0019
52796.4776	21	0.0018
52796.5402	22	0.0013
52796.7282	25	0.0001
52796.7889	26	-0.0022
52796.8531	27	-0.0011
52797.0425	30	-0.0009
52797.4186	36	-0.0031
52797.4800	37	-0.0048
52797.8025	42	0.0024
52798.1744	48	-0.0041
52798.2365	49	-0.0050
52798.2993	50	-0.0053
52798.4276	52	-0.0031
52798.4907	53	-0.0031
52798.6747	56	-0.0082
52798.9929	61	-0.0053
52799.1167	63	-0.0076
52799.1794	64	-0.0080
52799.2437	65	-0.0068
52799.3146	66	0.0011
52799.7486	73	-0.0063
52800.0104	77	0.0032
52800.0753	78	0.0051
52800.1928	80	-0.0036
52800.3299	82	0.0074
52801.2208	96	0.0154
52802.4025	115	-0.0010
52802.4651	116	-0.0014

References

Baba, H., Kato, T., Nogami, D., Hirata, R., Matsumoto, K., & Sadakane, K. 2000, PASJ, 52, 429
 Bailey, J. 1979, MNRAS, 189, 41P
 Belyavskij, A. I. 1933, Perem. Zvezdy, 4, 234
 Downes, R. A., & Margon, B. 1981, MNRAS, 197, 35P
 Hellier, C. 2001, Cataclysmic Variable Stars: how and why

Table 4. Fundamental characteristics of GO Com and LL And

	supercycle [†]	plateau [†]	amplitude
GO Com	~ 2800	8	4.8
LL And	~ 5000	9	~ 5

[†] unit in day

[‡] 1. This work 2. Howell, Hurst (1994), Kato (2004)

they vary (Berlin: Springer-Verlag)
 Howell, S. B., & Hurst, G. M. 1994, Inf. Bull. Variable Stars, 4043
 Howell, S. B., Szkody, P., & Cannizzo, J. K. 1995a, ApJ, 439, 337
 Howell, S. B., Szkody, P., Kreidl, T. J., Mason, K. O., & Puchnarewicz, E. M. 1990, PASP, 102, 758
 Howell, S. B., Szkody, P., Sonneborn, G., Fried, R., Mattei, J., Oliverson, R. J., Ingram, D., & Hurst, G. M. 1995b, ApJ, 453, 454
 Hu, J., Wei, J., Xu, D., Cao, L., & Dong, X. 1998, Ann. Shanghai Obs., Acad. Sin., 19, 235
 Ishioka, R., et al. 2003, PASJ, 55, 683
 Jurcevic, J. S., Honeycutt, R. K., Schlegel, E. M., & Webbink, R. F. 1994, PASP, 106, 481
 Kaluzny, J. 1986, IAU Circ., 4287
 Kato, T. 1997, PASJ, 49, 583
 Kato, T. 1999, Inf. Bull. Variable Stars, 4768
 Kato, T. 2004, PASJ, 56, S135
 Kato, T., & Hirata, R. 1990, Inf. Bull. Variable Stars, 3489
 Kato, T., Matsumoto, K., Nogami, D., Morikawa, K., & Kiyota, S. 2001a, PASJ, 53, 893
 Kato, T., Nogami, D., & Baba, H. 1995, Inf. Bull. Variable Stars, 4228
 Kato, T., Nogami, D., Moilanen, M., & Yamaoka, H. 2003a, PASJ, 55, 989
 Kato, T., et al. 2003b, MNRAS, 339, 861
 Kato, T., Sekine, Y., & Hirata, R. 2001b, PASJ, 53, 1191
 Kato, T., & Starkey, D. R. 2002, Inf. Bull. Variable Stars, 5358
 Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2004, PASJ, 56, S1
 Kowal, C. T. 1973, IAU Circ., 2562
 Lasota, J. P., Hameury, J. M., & Huré, J. M. 1995, A&A, 302, L29
 Marino, B. F., & Walker, W. S. G. 1979, in IAU Colloq. 46, Changing Trends in Variable Star Research, ed. F. M. Bateson, J. Smak, & J. H. Urch (Univ. of Waikato, Hamilton), N. Z., 29
 Mineshige, S., Liu, B., Meyer, F., & Meyer-Hofmeister, E. 1998, PASJ, 50, L5
 Moreau, O., & Reboul, H. 1995, A&AS, 111, 169
 Neustroev, V. V., Medvedev, A., Turbin, S., & Borisov, N. V. 2002, in ASP Conf. Ser. 261, The Physics of Cataclysmic Variables and Related Objects, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), 515
 Nogami, D., Baba, H., Kato, T., & Novák, R. 1998, PASJ, 50, 297
 Nogami, D., Masuda, S., & Kato, T. 1997, PASP, 109, 1114
 Nogami, D., Uemura, M., Ishioka, R., Kato, T., & Pietz, J. 2004, PASJ, 56, S155
 O'Donoghue, D., Chen, A., Marang, F., Mittaz, J. P. D., Winkler, H., & Warner, B. 1991, MNRAS, 250, 363
 Olech, A. 2003, Acta Astron., 53, 85




Fig. 1. The whole light curves of the 2003 superoutburst of GO Com. The zero point corresponds to magnitude 12.80, on a system close to R. On HJD 2452794, there shows a precursor with a decline rate of 0.74 mag d^{-1} .




Fig. 2. Theta diagram of a PDM period analysis of the data obtained during the plateau stage. The minimum point indicates 0.06297 d as the best-estimated superhump period.




Fig. 3. Phase-averaged light curve of GO Com superhumps, covering between HJD2452795 and HJD2452801, folded by 0.06359 d. The vertical and the horizontal axis denote the relative magnitude and the phase, respectively. A rapid rise and slower decline, which is a typical feature of superhumps, are shown.




Fig. 4. Phase-averaged light curves of GO Com superhumps. The abscissa and the ordinate are the phase of the superhump period and the differential magnitude, respectively.




Fig. 5. O-C diagram of superhump maxima. The abscissa and the ordinate denote the cycle count and $O - C$ (d) using equation (1), respectively. Typical error is 0.001 in each point. The quadratic fit corresponds to equation(2). There is a hint of a late superhump around HJD 2452802.




Fig. 6. Phase-averaged superhump profile on HJD2452794, corresponding to the precursor phase. There is no feature of superhump. This result was in contrast with the 1993 superoutburst of T Leo (Kato 1997) in which hump-like modulations, corresponding to the superhump period, was observed even in precursor.




Fig. 7. $P_{\text{dot}} = \dot{P}_{\text{SH}}/P_{\text{SH}}$ versus P_{SH} diagram. The data of GO Com are added to the original figure (figure 8 in Ishioka et al. 2003).

- Olech, A., Schwarzenberg-Czerny, A., Kędzierski, P.,
Złoczewski, K., Mularczyk, K., & Wiśniewski, M. 2003,
Acta Astron., 53, 175
- Osaki, Y. 1985, A&A, 144, 369
- Osaki, Y. 1995, PASJ, 47, 47
- Osaki, Y. 1996, PASP, 108, 39
- Osaki, Y., & Meyer, F. 2002, A&A, 383, 574
- Osaki, Y., & Meyer, F. 2003, A&A, 401, 325
- Patterson, J. 2001, PASP, 113, 736
- Patterson, J., Bond, H. E., Grauer, A. D., Shafter, A. W., &
Mattei, J. A. 1993, PASP, 105, 69
- Patterson, J., et al. 2002, PASP, 114, 721
- Ringwald, F. A., & Thorstensen, J. R. 1990, BAAS, 22, 1291
- Semeniuk, I., Olech, A., Kwast, T., & Nalezyty, M. 1997, Acta
Astron., 47, 201
- Stellingwerf, R. F. 1978, ApJ, 224, 953
- Uemura, M., et al. 2002a, PASJ, 54, 599
- Uemura, M., et al. 2002b, PASJ, 54, L15
- Usher, P. D. 1981, ApJS, 46, 117
- Verbunt, F., Bunk, W. H., Ritter, H., & Pfeffermann, E. 1997,
A&A, 327, 602
- Vogt, N., & Bateson, F. M. 1982, A&AS, 48, 383
- Warner, B. 1985, in Interacting Binaries, ed. P. P. Eggleton, &
J. E. Pringle (Dordrecht: D. Reidel Publishing Company),
367
- Warner, B. 1995, Cataclysmic Variable Stars (Cambridge:
Cambridge University Press)
- Wei, J., Jiang, X., Xu, D., Zhou, A., & Hu, J. 2001,
Chin.J.Astron.Astrophys, 1, 483
- Wenzel, W. 1982, Inf. Bull. Variable Stars, 2256

Date	HJD(start)*	HJD(end)	N [†]	Exp(s) [‡]	Observer
2003 June 3	52793.9623	52794.1228	170	60	TO
	52794.0083	52794.0213	27	30	GU
	52794.0407	52794.1488	461	10	ND
	52794.1546	52794.3686	254	60	TKr
2003 June 4	52795.0940	52795.1571	79	30	KYO
	52795.1515	52795.3608	141	120	TKr
2003 June 5	52795.6501	52795.7196	78	30	BY
	52795.7068	52795.8395	213	30	LC
	52795.9731	52796.0364	233	30	NH
	52795.9916	52796.1208	168	30	KYO
	52796.0051	52796.1106	467	10	ND
	52796.0360	52796.1177	85	60	TA
	52796.0408	52796.0843	499	3	SY
	52796.1519	52796.2097	25	120	TKr
	52796.3758	52796.5475	270	40	TV
	52796.6002	52796.7288	94	90	DS
2003 June 6	52796.6675	52796.8522	513	30	BY
	52796.7079	52796.8725	262	30	LC
	52796.9587	52797.0507	100	30	NH
	52796.9613	52797.1570	87	60	TO
	52797.0199	52797.0878	408	3	SY
	52797.1044	52797.2231	24	30	KYO
	52797.3882	52797.5218	150	30	JP
	52797.4377	52797.5400	120	60	TV
	52797.7359	52797.8367	57	120	MA
	52798.1500	52798.3598	115	120	TKr
2003 June 7	52798.3781	52798.5449	273	40	TV
	52798.3858	52798.5346	173	30	JP
	52798.6178	52798.7296	85	90	DS
	52798.6947	52798.7165	15	30	LC
2003 June 8	52798.9797	52799.1420	189	30	KYO
	52798.9825	52799.1028	442	10	ND
	52798.9985	52799.1212	434	20	SY
	52799.0367	52799.1069	80	60	TA
	52799.1528	52799.3560	139	60	TKr
	52799.4082	52799.4738	102	40	TV
	52799.6882	52799.7583	157	30	LC
	52799.9696	52800.0387	419	20	SY
2003 June 9	52800.1506	52800.3421	132	60	TKr
	52800.3789	52800.4222	40	45	JP
	52800.5842	52800.7114	104	90	DS
	52800.9717	52801.0623	273	20	SY
2003 June 10	52801.1504	52801.2829	86	60	TKr
	52802.0140	52802.1140	92	20	SY
	52802.1569	52802.3072	61	60	TKr
	52802.3782	52802.5009	126	45	JP
2003 June 11	52802.3848	52802.5333	203	50	TV
	52802.7268	52802.8500	75	120	MA
	52803.1551	52803.3364	122	60	TKr
	52803.9766	52804.0338	62	30	NH
2003 June 12	52804.1516	52804.3296	123	60	TKr
	52804.3974	52804.5200	167	50	TV
	52805.1537	52805.2685	78	60	TKr
2003 June 13	52805.5862	52805.7097	105	90	DS
	52805.9736	52806.0933	246	20	SY
	52806.1573	52806.3333	107	120	TKr
2003 June 14	52806.5884	52806.7076	92	90	DS

Table 1. (Continued)

	52807.1613	52807.3274	58	120	TKr
2003 June 17	52808.1572	52808.3105	54	240	TKr
2003 June 18	52808.6183	52808.7090	71	90	DS
2003 June 19	52810.6007	52810.6362	31	90	DS

* HJD-2400000

† Number of frames

‡ Exposure times

Instrument TO:25cm+Apogee AP-7(Saitama, Japan),

GU:25cm+AP7p (Gunma, Japan),

ND:60cm+PixCellentS/T 00-3194(SITe 003AB)(Hida, Japan)

KYO:30cm+ST-7E(Kyoto, Japan)

LC:73cm+GenesisG16+90KAF1602E(California, USA)

NH:25cm+CV-04(Mie, Japan)

TA:21cm+ST-7(Okayama, Japan)

SY:28cm+ST-9E(Nayoro, Japan)

TV:35cm+ST-7 (Landen, Belgium)

DS:36cm+ST-10XME(Indiana, USA)

BY:45cm+PRIMEFOCUS (Arizona, USA)

JP:20cm+ST-6B (Erfstadt, Germany)

MA:12.5cm+ST-7E (Alberta, Canada)

TKr:28cm+ST-7E (Tashkent, Uzbekistan)

Table 2. Recorded outbursts of GO Com

	day	JD	type*	references†	
Jul. 1973	2441864		?		1
May. 1989	2447676		S?		2
Apr. 1990	2448011		N?		3
Jul. 1995	2449914		S		4
Mar. 1996	2450168		N		2
Feb. 1997	2450486		N		2
Apr. 1998	2450558		N		2
Jun. 2003	2452794		S		5

* S: superoutburst, N: normal outburst

† 1. Kowal 1973 2. VSNET archives 3. Kato, Hirata 1990 4. Kato et al. 1995 5. this work